

Utilization of a Next Generation Decellularized Dermal Allograft to Augment a Recurrent Peroneus Brevis Tear

Vytautas Ringus, MD

Oklahoma City, OK

Board Certified, American Board of Orthopaedic Surgeons

Surgical Summary

Clinical Presentation:

- 54-year-old female with 2 previous peroneal tendon repairs presented with sharp lateral pain over the peroneal tendon.
- Exam revealed increased pain with resisted inversion and eversion, local swelling, 4/5 strength on testing, no restriction in range of motion, and no subluxation of tendons.
- MRI revealed significant scarring and a recurrent peroneus brevis tear (Figure 1).

Intraoperative Findings:

- Exploration revealed a large retrofibular tear of the brevis tendon, complex delaminating intrasubstance tear of the peroneal tendons with splitting of the brevis, and tenosynovitis of both the peroneus longus and brevis.
- The tear extended proximal to the fibula, and distal to the peroneal tubercle.




Figure 1

Surgeon Perspective:

"DermaPure provided reinforcement of this repair as well as protection from mechanical stress. This patient had reduced scarring."

Utilization of a Next Generation Decellularized Dermal Allograft to Augment a Recurrent Peroneus Brevis Tear

Vytautas Ringus, MD

Oklahoma City, OK

Board Certified, American Board of Orthopaedic Surgeons

Surgical Procedure:

- A lateral approach was taken utilizing her previous incision. The superior peroneal retinaculum was incised, and both tendons visualized.
- Debridement and repair of the large retrofibular tear of the brevis, and debridement of the longus.
- DermaPure[®] decellularized dermal allograft was applied in a tubularized fashion around the brevis, (due to the history of multiple procedures, extent of scarring, and integrity of the tendon), to augment and stimulate healing, while minimizing scarring to improve gliding kinematics (Figure 2).
- The retrofibular groove was palpated and noted to be appropriately deep, and the superior peroneal retinaculum was imbricated with absorbable suture.




Figure 2

DermaPure[®] Implantation:

- 3 cm x 4 cm
- Basement membrane outer-most

Post-Operative Note:

- Patient was discharged home, non-weight bearing and immobilized for 6 weeks.
- At 6 weeks, patient was placed in a boot and physical therapy began.
- Patient is being followed routinely to assess clinical outcomes.

TissueRegenixUS.com

833-567-0767

1808 Universal City Blvd, Universal City, TX 78148